

# HW#4

Due 2023/11/20

1. Show that

$$\rho \frac{d\vec{u}}{dt} = -\nabla P + \rho_c \vec{E} + \vec{J} \times \vec{B}$$

can be rewritten as

$$\frac{\partial}{\partial t} \left( \rho \vec{u} + \frac{1}{C^2} \frac{\vec{E} \times \vec{B}}{\mu_0} \right) + \nabla \cdot \left( \rho \vec{u} \vec{u} + \left( P + \frac{1}{2} \epsilon_0 E^2 + \frac{B^2}{2\mu_0} \right) \vec{1} - \left( \epsilon_0 \vec{E} \vec{E} + \frac{\vec{B} \vec{B}}{\mu_0} \right) \right) = 0$$

First, you need to show that  $\frac{\partial}{\partial t} (\rho \vec{u}) + \nabla \cdot (\rho \vec{u} \vec{u}) = \rho \frac{d\vec{u}}{dt}$

by using  $\nabla \cdot (\vec{A} \vec{B}) = (\nabla \cdot \vec{A}) \vec{B} + (\vec{A} \cdot \nabla) \vec{B}$

2. Show that

$$\frac{\partial}{\partial t} \left( \frac{1}{2} \rho u^2 + U_{int} \right) + \nabla \cdot \left( \left( \frac{1}{2} \rho u^2 + U_{int} \right) \vec{u} + P\vec{u} + Q \right) = \vec{J} \cdot \vec{E}$$

can be rewritten as

$$\frac{\partial}{\partial t} \left( \frac{1}{2} \rho u^2 + U_{int} + \frac{1}{2} \epsilon_0 E^2 + \frac{B^2}{2\mu_0} \right) + \nabla \cdot \left( \left( \frac{1}{2} \rho u^2 + U_{int} \right) \vec{u} + P\vec{u} + Q + \frac{\vec{E} \times \vec{B}}{\mu_0} \right) = 0$$